

TERNARY QUARTICS AND 3-DIMENSIONAL COMMUTATIVE ALGEBRAS

P.KATSYLO, D.MIKHAILOV

May 5, 1994

ABSTRACT. We find the connection between 3-dimensional commutative algebras with a trivial trace and plane quartics and its bitangents.

§0. A structure of a commutative algebra on $\mathbb{C}^3$ is called 3-dimensional algebra. Let $\mathcal{A}$ be the set of 3-dimensional algebras. Consider $\mathcal{A}$ as a linear space. Let $\mathcal{A}_0 \subset \mathcal{A}$ be the linear subspace of algebras with a trivial trace. By definition, $\eta \in \mathcal{A}_0$ iff the contraction of the structural constants of η is equal to zero.

By PV denote the projectivization of a vector space V . For $v \in V$, $\bar{v} \neq 0$ denote by $\bar{v}$ the corresponding point of the projective space PV .

Let $\eta \in \mathcal{A}_0$ be an algebra with a trivial trace. Recall that an element $a \in \mathbb{C}^3$ is called an idempotent, iff $a \neq 0$, $a^2 = a$. We say that an element $\bar{a} \in P\mathbb{C}^3 = \mathbb{P}^2$ is called a generalized idempotent, iff $a^2 = \lambda a$, there $\lambda \in \mathbb{C}$. Every idempotent defines a generalized idempotent. Every generalized idempotent $\bar{a} \in \mathbb{P}^2$ such that $a^2 \neq 0$ defines uniquely the idempotent $a' \in \mathbb{C}^3$ such that $\bar{a} = \overline{a'}$. Define the subscheme $X(\eta) \subset \mathbb{P}^2$ of the generalized idempotents by the following equation:

$$(0.1) \quad a^2 \wedge a = 0$$

Consider the open SL_3 -invariant subset

$$\mathcal{A}'_0 = \{\eta \in \mathcal{A}_0 \mid \dim X(\eta) = 0\} \subset \mathcal{A}_0.$$

Lemma 0.1. $\mathcal{A}'_0$ is nonempty

Consider $\eta \in \mathcal{A}_0$. The algebra η defines the quadratic mapping

$$\mathbb{C}^3 \longrightarrow \mathbb{C}^3, \quad a \mapsto a^2.$$

This quadratic mapping defines the section $\tilde{\eta}$ of the vector bundle $T_{\mathbb{P}^2}(1)$. The scheme of zeros of the section $\tilde{\eta}$ is $X(\eta)$. We have

$$\deg X(\eta) = c_2(T_{\mathbb{P}^2}(1)) = 7,$$

Research supported by Max-Planck-Institute fur Mathematik and Grant N MQZ000 of the International Science Foundation.

for $\eta \in \mathcal{A}'_0$.

Consider the open SL_3 -invariant subset

$$\begin{aligned} \mathcal{A}''_0 = \{ & \eta \in \mathcal{A}'_0 \mid X(\eta) = \{\overline{a_1}, \dots, \overline{a_7}\}, a_i^2 \neq 0, 1 \leq i \leq 7, \\ & \text{every 3 points of } X(\eta) \text{ does not lie on a line,} \\ & \text{every 6 points of } X(\eta) \text{ does not lie on a quadric} \} \subset \mathcal{A}'_0. \end{aligned}$$

Lemma 0.2. $\mathcal{A}''_0$ is nonempty.

Consider the rational SL_3 -morphism

$$\phi : P\mathcal{A}_0 \longrightarrow (\mathbb{P}^2)^{(7)}, \quad \eta \mapsto X(\eta).$$

Proposition 0.3. ϕ is a birational isomorphism.

In other words, 3-dimensional algebra in general position with a trivial trace is uniquely (up to a scalar factor) defined by its generalized idempotents.

Fix $\eta \in \mathcal{A}''_0$. Let $a_1, \dots, a_7$ be the idempotents of the algebra η . Let

$$\pi = \pi(\eta) : Z = Z(\eta) \longrightarrow \mathbb{P}^2$$

be the blowing up of $X(\eta)$ in $\mathbb{P}^2$. It is well known that Z is Del Pezzo surface of degree 2. Let

$$\beta = \beta(\eta) : Z \longrightarrow \mathbb{P}^{2*}$$

be the canonical double covering with the nonsingular quartic $Y = Y(\eta) \subset \mathbb{P}^{2*}$ as the branch locus.

SL_3 -module $S^2\mathcal{A}_0$ contains with multiplicity 1 the isomorphic to $S^4\mathbb{C}^3$ submodule. Therefore, there exists a unique (up to a scalar factor) nontrivial quadratic SL_3 -mapping

$$\epsilon : \mathcal{A}_0 \longrightarrow S^4\mathbb{C}^3.$$

Lemma 0.4. $\epsilon(\eta) \neq 0$.

Consider the quaternary form $\epsilon(\eta)$ on the space $\mathbb{C}^{3*}$. This quaternary form defines quartic $Y' = Y'(\eta) \subset \mathbb{P}^{2*}$. Consider 28 linear forms $a_1, \dots, a_7, (a_i - a_j)^2, 1 \leq i < j \leq 7$ on the space $\mathbb{C}^{3*}$. These linear forms define 28 lines $A_1, \dots, A_7, A_{ij} \in \mathbb{P}^{2*}$.

Theorem 0.5. 1) Y is isomorphic to Y' .

2) 28 bitangents to Y' are $A_1, \dots, A_7, A_{ij}, 1 \leq i < j \leq 7$.

The authors thank B.Broer for useful discussion.

§1. Let e_1, e_2, e_3 be the standart basis in $\mathbb{C}^3$, and x_1, x_2, x_3 be the dual basis in $\mathbb{C}^{3*}$. The group SL_3 acts canonically in the space $S^a\mathbb{C}^3 \otimes S^b\mathbb{C}^{3*}$, $a, b \geq 0$. For $a, b \geq 1$ consider the linear SL_3 -mapping

$$\Delta = \sum \frac{\partial}{\partial e_i} \otimes \frac{\partial}{\partial x_i} : S^a\mathbb{C}^3 \otimes S^b\mathbb{C}^{3*} \longrightarrow S^{a-1}\mathbb{C}^3 \otimes S^{b-1}\mathbb{C}^{3*}.$$

It is well known that a representation of the group SL_3 in the space $V(a, b) = \ker \delta$ is irreducible. Assume that $V(a, 0) = S^a\mathbb{C}^3, V(0, b) = S^b\mathbb{C}^{3*}$.

A structure of a commutative algebra on $\mathbb{C}^3$ is a symmetric bilinear mapping

$$\mathbb{C}^3 \times \mathbb{C}^3 \longrightarrow \mathbb{C}^3.$$

The set of such symmetric bilinear mappings is $\mathbb{C}^3 \otimes S^2\mathbb{C}^{3*}$. Therefore, the linear space $\mathcal{A}$ of 3-dimensional algebras is $\mathbb{C}^3 \otimes S^2\mathbb{C}^{3*}$. The contraction of structural constants of algebras is the mapping

$$\Delta : \mathbb{C}^3 \otimes S^2\mathbb{C}^{3*} \longrightarrow \mathbb{C}^{3*}.$$

Therefore, the linear space $\mathcal{A}_0$ of 3-dimensional algebras with a trivial trace is $V(1, 2)$.

The decomposition

$$V(1, 2) \otimes \mathbb{C}^3 \otimes \mathbb{C}^3 \simeq \mathbb{C}^3 \oplus 2V(0, 2) \oplus 2V(2, 1) \oplus 2V(1, 3) \oplus V(3, 2)$$

holds. Thus there exists a unique (up to a scalar factor) nontrivial trilinear SL_3 -mapping

$$\mu : V(1, 2) \times \mathbb{C}^3 \times \mathbb{C}^3 \longrightarrow \mathbb{C}^3.$$

Let us give an explicite form of μ :

$$\mu(e_{i_1} \otimes x_{j_1} x_{j_2}, e_{i_2} e_{i_3}) = \Delta^2(e_{i_1} e_{i_2} e_{i_3} \otimes x_{j_1} x_{j_2}).$$

The corresponding to $\eta \in V(1, 2)$ algebraic structure is the bilinear symmetric mapping

$$\mu(\eta, \cdot, \cdot) : \mathbb{C}^3 \times \mathbb{C}^3 \longrightarrow \mathbb{C}^3.$$

Example 1.1. Consider the algebra

$$\eta_0 = \frac{1}{4}(e_1 \otimes x_3^2 + e_2 \otimes x_1^2 + e_3 \otimes x_2^2).$$

The multiplication table of η_0 is as following:

$$e_1 * e_2 = e_2 * e_3 = e_3 * e_1 = 0, \quad e_1^2 = e_2, \quad e_2^2 = e_3, \quad e_3^2 = e_1.$$

It can be easily checked that the subscheme $X(\eta_0) \subset \mathbb{P}^2$ of the generalized idempotents of η_0 is

$$X(\eta_0) = \{\overline{\theta e_1 + \theta^2 e_2 + \theta^4 e_3} \mid \theta \in \mu_7\}.$$

Proof of lemmas 0.1 and 0.2. We have $\eta_0 \in \mathcal{A}'_0$. It follows that $\mathcal{A}'_0$ is nonempty. It can easily be checked that $\eta_0 \in \mathcal{A}''_0$. Therefore, $\mathcal{A}''_0$ is nonempty.

Proof of proposition 0.3. It can easily be checked that $\phi^{-1}(X(\eta_0)) = \overline{\eta_0}$. It follows from (0.1) that a closure of a fiber of ϕ is a linear subspace in $PV(1, 2)$ of dimension $\leq \dim PV(1, 2) - \dim(\mathbb{P}^2)^{(7)} = 0$. Hence ϕ is a birational isomorphism.

§2. Fix $\eta \in \mathcal{A}''_0$. Let $a_1, \dots, a_7$ be the idempotents of η .

Consider the cubic mapping

$$\psi = \psi(\eta) : \mathbb{C}^3 \longrightarrow \wedge^2 \mathbb{C}^3 \simeq \mathbb{C}^{3*}, \quad a \mapsto a^2 \wedge a.$$

Lemma 2.1. *Consider a_i as a linear form on $\mathbb{C}^{3*}$. Let $Q_i = Q_i(\eta)$ be the corresponding to the cubic form $\psi^*(a_i)$ cubic. Then the cubic Q_i contains $\overline{a_1}, \dots, \overline{a_7}$. Moreover, Q_i contains $\overline{a_i}$ with multiplicity ≥ 2 .*

Proof. It is obvious that $Q_i \ni \overline{a_1}, \dots, \overline{a_7}$. Let us prove that Q_i contains $\overline{a_i}$ with multiplicity ≥ 2 . We have

$$\begin{aligned} \psi^*(a_i) : a &\mapsto a^2 \wedge a \wedge a_i \in \wedge^3 \mathbb{C}^3 \simeq \mathbb{C}, \\ \psi^*(a_i)(a_i + tb) &= (a_i + tb)^2 \wedge (a_i + tb) \wedge a_i = a_i^2 \wedge a_i \wedge a_i + \\ &+ t((2a_i * b) \wedge a_i \wedge a_i + a_i^2 \wedge b \wedge a_i) + \dots = 0 + t \cdot 0 + \dots \end{aligned}$$

for any $b \in \mathbb{C}^3$.

Lemma 2.2. *Consider $(a_i - a_j)^2, i < j$ as a linear form on $\mathbb{C}^{3*}$. Let $Q_{ij} = Q_{ij}(\eta)$ be the corresponding to a cubic form $\psi^*((a_i - a_j)^2)$ cubic. Then Q_{ij} is the union of the line $\langle \overline{a_i}, \overline{a_j} \rangle$ and the containing the points $\overline{a_1}, \dots, \widehat{\overline{a_i}}, \dots, \widehat{\overline{a_j}}, \dots, \overline{a_7}$ quadric.*

Proof. It is obvious that $Q_{ij} \ni \overline{a_1}, \dots, \overline{a_7}$. We have to prove that Q_{ij} contains the line $\langle \overline{a_i}, \overline{a_j} \rangle$. We have

$$\begin{aligned} \psi^*((a_i - a_j)^2) : a &\mapsto a^2 \wedge a \wedge (a_i - a_j)^2 \in \wedge^3 \mathbb{C}^3 \simeq \mathbb{C}, \\ \psi^*((a_i - a_j)^2)(t_i a_i + t_j a_j) &= (t_i a_i + t_j a_j)^2 \wedge (t_i a_i + t_j a_j) \wedge (a_i - 2a_i * a_j + a_j) = \\ &= t_i^3 a_i^2 \wedge a_i \wedge (a_i - 2a_i * a_j + a_j) + t_i^2 t_j (a_i^2 \wedge a_j \wedge (a_i - 2a_i * a_j + a_j) + \\ &+ ((2a_i * a_j) \wedge a_i \wedge (a_i - 2a_i * a_j + a_j))) + \dots = t_i^3 \cdot 0 + t_i^2 t_j \cdot 0 + \dots \equiv 0. \end{aligned}$$

Consider the rational morphism

$$\Psi = \Psi(\eta) : \mathbb{P}^2 \longrightarrow \mathbb{P}^{2*}, \quad \overline{a} \mapsto \overline{\psi(a)}.$$

The rational morphism Ψ is not defined on $X(\eta)$. Let

$$\mathbb{P}^2 \xleftarrow{\pi=\pi(\eta)} Z = Z(\eta) \xrightarrow{\beta=\beta(\eta)} \mathbb{P}^{2*}$$

be the regularization of Ψ . It is well known that Z is Del Pezzo surface of degree 2, π is the blowing up of the seven points $\overline{a_1}, \dots, \overline{a_7}$ in $\mathbb{P}^2$, $\beta : Z \longrightarrow \mathbb{P}^{2*}$ is double covering with the nonsingular quartic $Y = Y(\eta) \subset \mathbb{P}^{2*}$ as the branch locus.

Proof of lemma 0.4 and theorem 0.5. Consider the nontrivial homogeneous (of degree 6) SL_3 -mappings

$$\begin{aligned} \gamma_i : V(1, 2) &\longrightarrow V(0, 12) = S^{12} \mathbb{C}^{3*}, \\ \gamma_1 : \eta &\mapsto \psi^*(\epsilon(\eta)), \\ \gamma_2 : \eta &\mapsto \left(\det \left(\frac{\partial \psi^*(e_i)}{\partial x_j} \right) \right)^2. \end{aligned}$$

The decomposition

$$S^6 V(1, 2) \simeq \dots \oplus V(1, 12) \oplus \dots$$

holds. Thus $\gamma_1 = c\gamma_2$, there $c \neq 0$. This implies lemma 0.4 and statement 1) of theorem 0.5.

Statement 2) of theorem 0.5 is a corollary of lemmas 2.1 and 2.2 (see [1])

REFERENCES

1. Ph. Griffiths and J. Harris, *Principles of algebraic geometry*, Wiley-Interscience, 1978.